

Fundamental Markov systems

Ivan Werner

Moscow, Russia

Email: ivan_werner@mail.ru

February 9, 2009

Abstract

We continue development of the theory of Markov systems initiated in [34]. In this paper, we introduce fundamental Markov systems associated with random dynamical systems and show that the proof of the uniqueness and empiricalness of the stationary initial distribution of the random dynamical system reduces to that for the fundamental Markov system associated with it. The stability criteria for the latter are much clearer.

MSC: 60J05, 37A50, 37H99, 28A80

Keywords: Markov systems, random dynamical systems, iterated function systems with place-dependent probabilities, random systems with complete connections, g -measures, Markov chains, fractals.

In [34], the author initiated the study of a general concept of a Markov system. This was motivated by a desire to have a scientifically consistent unifying mathematical structure which would cover *finite Markov chains* [10], g -measures [25] and *iterated function systems with place-dependent probabilities* [1], [13].

The purpose of this note is to show that the structure of a Markov system arises naturally (possibly unavoidably) in the study of *random dynamical systems*.

1 Random dynamical systems

Let (K, d) be a complete separable metric space and E a countable set. For each $e \in E$ let a Borel measurable map $w_e : K \rightarrow K$ and a Borel measurable probability function $p_e : K \rightarrow [0, 1]$ be given, i.e.

$$\sum_{e \in E} p_e(x) = 1 \quad \text{for all } x \in K.$$

We call the family $\mathcal{D} := (K, w_e, p_e)_{e \in E}$ a *random dynamical system*. A survey on random dynamical systems can be found e.g. in [23]. If a reader doesn't see, how the definition of random dynamical system in [23] relates to that in this paper, it is explained in [24].

With the random dynamical system $\mathcal{D}$ is associated a *Markov operator* U acting on all bounded Borel measurable functions f by

$$Uf := \sum_{e \in E} p_e f \circ w_e$$

and its *adjoint operator* acting on the set of Borel probability measures ν by

$$U^*\nu(f) := \int U(f) d\nu.$$

A measure μ is called *invariant* with respect to the random dynamical systems if and only if

$$U^*\mu = \mu.$$

Let $\Sigma^+ := \{(\sigma_1, \sigma_2, \dots) : \sigma_i \in E, i \in \mathbb{N}\}$ endowed with the product topology of discrete topologies and $S : \Sigma^+ \rightarrow \Sigma^+$ be the left shift map. For $x \in K$, let P_x be the Borel probability measure on Σ^+ given by

$$P_x([e_1, \dots, e_n]) := p_{e_1}(x)p_{e_2} \circ w_{e_1}(x) \dots p_{e_n} \circ w_{e_{n-1}} \circ \dots \circ w_{e_1}(x),$$

for every cylinder set $[e_1, \dots, e_n] := \{\sigma \in \Sigma^+ : \sigma_1 = e_1, \dots, \sigma_n = e_n\}$, which is associated with the Markov process generated by the random dynamical system with the Dirac initial distribution δ_x .

Remark 1 Note that each map w_e needs to be defined only on a subset of K where its probability function p_e is greater than zero. In this case, one obtains a random dynamical system on K by extending the maps on the whole space arbitrarily.

1.1 Historical context

First, I will give some historical roots of the theory, and then, I will list some works which in my view form the historical context of this paper.

It was not clear until the beginning of the twentieth century whether the independence of random variables is a necessary condition for the *law of large numbers* and the *central limit theorem* to hold. The breakthrough was a work by A. A. Markov [26] in which he has extended the *law of large numbers* to dependent random variables (he also extended the central limit theorem to such processes, see [11] for a nice account on Markov's work and life). Markov restricted himself to processes where each random variable depends only on the previous one. Such a process, in case of a discrete state space, is generated by a transition matrix (or a directed graph with probability weights) and an initial distribution. These processes, now known as *Markov chains*, found many applications.

From the work of Markov naturally arises the question whether the law of large numbers and other limit theorems hold true for a more general class of dependent processes. The truly next class of processes can be only those where the dependence of a random variable on the past is not restricted to any number of previous variables. The study of such processes was initiated (and motivated by applications) by O. Onicescu and G. Mihoc [27]. Remarkably, they found a way of constructing such processes without giving infinitely many rules of dependence of a random variable on the values of all previous, which is of course very important for applications. They called such processes *les chaînes à liaisons complètes*. Their work gave rise to the theory of *dependence with complete connections*, in which many limit theorems have been proved [17]. What I call in this paper *random dynamical system* can be seen as a special case of this theory. However, it must be said that every stationary process with values in E can already be generated by such a random dynamical system. Let's illustrate the dependence in the notation of this paper. If maps w_e are contractions on a complete metric space, the past is coded by the maps to a point in topological space K and the probability of a value from E of the next random variable is then obtained as a function evaluated at that point,

$$P(X_1 = e_1 | X_0 = e_0, X_{-1} = e_{-1}, \dots) = p_{e_1} \left(\lim_{n \rightarrow -\infty} w_{e_0} \circ w_{e_{-1}} \circ \dots \circ w_{e_{-n}}(x_0) \right)$$

for all $(\dots, e_{-1}, e_0, e_1, \dots) \in E^{\mathbb{Z}}$ and $x_0 \in K$.

One could argue that the development so far was guided mainly by the internal mathematical logic. However, the main reason for the development of the mathematical language still is the striving of Homo Sapiens for a better description of the world outside (even if many modern mathematical craftsmen have no interest in science at all). The random dynamical systems such as in this paper arose naturally also as *learning models* [20], [9]. Consider an intelligent object, e.g. a rat. Its state of "intelligence" x is assumed to take some value from $[0, 1] \subset \mathbb{R}$. The object is "asked some questions" and its responses are measured as '0' ("wrong") or '1' ("right"). If it gives an answer '0', its "intelligence" is moved to $w_0(x) := 1/2x$. Otherwise, its "intelligence" is moved to $w_1(x) := 1/2x + 1/2$. It is natural to assume that a response '0' can happen with some probability $p_0(x)$ (depending on the current state of "intelligence") and a response '1' with probability $p_1(x) := 1 - p_0(x)$. One assumes that the function p_0 does not change with the time (if it did, it's likely that it would have some time average, which only would matter [3]). Furthermore, probability functions p_0 and p_1 easily can be obtained empirically if sufficiently many measurements are made. Many seminal works on Markov processes generated by random dynamical systems have been motivated by learning models.

The striving for a classification of stationary random processes brought attention to such systems once again. After Ornstein [28] had shown that Kolmogorov-Sinai entropy is a complete invariant on Bernoulli shifts and not complete on Kolmogorov automorphisms [29], arose an interest in measure preserving trans-

formations which are Kolmogorov and not Bernoulli. Of course, a natural candidate for that is a strongly mixing process on a finite alphabet with infinite memory, which is known to be generated by these kind of random dynamical systems. It was M. Keane [21] who drew attention of ergodic theory people to such systems. He considered a special case where the maps of the system can be obtained as inverse branches of a (expanding) homomorphism S of a compact metric space K (e.g. $K := E^{\mathbb{N} \cup 0}$ and S is shift map, then $w_e(\dots, y_{-1}, y_0) := (\dots, y_{-1}, y_0, e)$ for all $(\dots, y_{-1}, y_0) \in K$). In this case, the probabilities for these maps can be given in terms of a function $g : K \rightarrow [0, 1]$ ($p_e(\dots, y_{-1}, y_0) := g(\dots, y_{-1}, y_0, e)$ for all $(\dots, y_{-1}, y_0) \in K$) with the property that $\sum_{y \in S^{-1}(\{x\})} g(y) = 1$ for all $x \in K$. Then the stationary process also is invariant with respect to homomorphism S . He called the stationary process in this case a *g-measure*. Much of the progress on the subject has been made from the study of this special case. Of course, the one-sided symbolic space is more than just an example. However, the restriction to this case clearly encourages an algebraic rather than a geometric approach, which could be one of the reasons why the structure of *the fundamental Markov system*, which is going to be introduced in this paper, can not have been seen before.

A careful reader probably has noticed that in all of the examples so far the maps of the system have been contractive. It was shown by J. Hutchinson [16] that a family of contractive maps on a compact metric space has a unique invariant subset. These subsets are like footprints of this kind of "animals". They are often very irregular on every scale. (This must be very fascinating already for itself because there are even people who study these footprints without actually any interest in any "animal".) Advances in computational technology allowed to produce beautiful colorful pictures of such sets. Naturally, arose the idea of storing images with such systems [14]. Moreover, contractive maps allow to code strings of symbols such as '0' and '1' to a point in the state space. This can be useful for data compression [4]. This all attracted a new interest to the random dynamical systems. The study of such systems for these purposes was continued mainly by M. F. Barnsley *et al.* [1], [2]. They called them *iterated function systems with place dependent probabilities*. Much of the literature on the subject is now available also under this name.

Recently, such system have been suggested also as a natural model for a quantum measurement process [31]. Let K be a state space of a quantum system and E a set of possible outputs of a measurement apparatus. When the quantum system being in a state $x \in K$ interacts with the measurement apparatus and we observe $e \in E$, it is known that this usually results in the system moving to a new state $y \in K$. Hence, it is natural to assume that there is a map $w_e : K \rightarrow K$ associated with each output $e \in E$. Moreover, it is natural to assume that e could be observed with some probability $p_e(x)$ in this experiment. This all also can be formulated using the conventional language of Hilbert space and projection operators [31]. The reader probably has noticed the analogy with the setup of the "learning model" above. It probably is an indication on the universality of the approach. In fact, studying a system by asking it some

"question" is a very natural approach.

I hope I have convinced the reader that the structure which is being studied in this paper is not one of the toys designed by some mathematicians to keep them busy, but one which naturally has been crystalizing in the modern science. It clearly needs to be properly integrated into the body of mathematics which describes deterministic, random and quantum paradigms. Now, I will list some important contributions which were made for the purpose of understanding such systems with respect to their stability and ergodic properties which clearly form the historical context of this paper.

W. Doeblin and R. Fortet [12] (1937) gave fairly weak condition on strictly positive probability functions on a compact metric space which insures that the system with contractive maps has a unique (attractive) stationary initial distribution. In particular, this condition is satisfied if the probabilities have a summable variation (Dini-continuous).

L. Breiman [9] (1960) proved the strong law of large numbers for Markov operators with the Feller property which posses a unique stationary initial distribution on a compact Hausdorff space (see a sharper result in [37]).

R. Isaac [18] (1962) introduced the average contractiveness condition, which insures the uniqueness of the stationary state (he proved it on a compact metric space with strictly positive probability functions satisfying Lipschitz continuity).

F. Ledrappier [25] (1974) identified the g -measures as projections of some equilibrium states defined by variational principle with respect to the potential $\log g$ seen as a function on $E^{\mathbb{Z}}$ (see [38] for the explanation in the general case). Furthermore, he showed that the natural extension of the g -measure is weakly Bernoulli if the g -function is strictly positive and Dini-continuous. P. Walters [33] (1975) extended the result to such g on a subshift of finite type (Markov system).

T. Kaijser [22] (1981) introduced a local contractiveness on average condition in the general setup of random systems with complete connections, which also can imply the uniqueness of the stationary state. He called his systems *weakly distance diminishing random systems with complete connections*.

H. Berbee [5] (1987) showed the uniqueness and the very weak Bernoulli property of the g -measure for strictly positive g -functions on a full shift satisfying a continuity condition which is weaker than the Dini-continuity. After that, many other works have been devoted to the weakening of the algebraic expression forcing a continuity of the g -function, but mostly only for the proof of uniqueness of the stationary state (see Ö. Stenflo [30] (2003) and N. Berger, Ch. Hoffman, V. Sidoravicius [6] (2005) and the references there). (It must be pointed out that the continuity of the probability functions is not fundamental for the stability of such systems (e.g. Example 2)).

J. H. Elton [13] (1987) recognized the importance of the relation for $x, y \in K$ given by the equivalence of measures P_x and P_y for the proof of the ergodic

theorem for such systems (though, he still assumed the uniqueness of the stationary state, which was shown later [35] to be not necessary). He proved that this equivalence relation holds true for all $x, y \in K$ if all probability functions are bounded away from zero, Dini-continuous and the system satisfies a contractiveness on average condition (in the language of this paper, the fundamental Markov system associated with such a random dynamical system has a single vertex set).

M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo [1] [2] (1989) made, in my view, two important contributions to understanding the conditions for the stability of such systems. They showed that the condition of the strict positivity of the probability functions can be weakened (is not fundamental). Secondly, they found implicitly a way of reduction of the multiplicative average contractiveness condition to the additive average contractiveness condition, though they did not accomplish it completely (see [34] for details).

I. Werner [35] (2005) showed that the condition of equivalence of measures P_x and P_y for all x and y in the same vertex set of a Markov system (see next section), which is continuous, irreducible and contractive, is sufficient for the uniqueness of the stationary initial distribution. For example (the example was given by using some ideas of A. Johansson and A. Öberg [19]), this condition is satisfied if the probability functions have a square summable variation on each vertex set and are bounded away from zero, which is more general than Elton's [13] example.

This list is far from being complete. There are many other works, which a reader can easily find under code names *g-measures*, *iterated functions systems with place-dependent probabilities*, *random systems with complete connections*, *random dynamical systems* and *Markov systems*. A reader interested in the study of general Markov operators is referred to [32].

1.2 Markov systems

Now, let us consider a special random dynamical system which we call a *Markov system* [34].

Let $K_1, K_2, \dots, K_N$ be a partition of a metric space K into non-empty Borel subsets (we do not exclude the case $N = 1$). Furthermore, for each $i \in \{1, 2, \dots, N\}$, let

$$w_{i1}, w_{i2}, \dots, w_{iL_i} : K_i \longrightarrow K$$

be a family of Borel measurable maps such that for each $j \in \{1, 2, \dots, L_i\}$ there exists $n \in \{1, 2, \dots, N\}$ such that $w_{ij}(K_i) \subset K_n$ (Fig. 1). Finally, for each $i \in \{1, 2, \dots, N\}$, let

$$p_{i1}, p_{i2}, \dots, p_{iL_i} : K_i \longrightarrow \mathbb{R}^+$$

be a family of positive Borel measurable probability functions (associated with the maps), i.e. $p_{ij} > 0$ for all j and $\sum_{j=1}^{L_i} p_{ij}(x) = 1$ for all $x \in K_i$.




Fig. 1. A Markov system.

Definition 1 We call $V := \{1, \dots, N\}$ the set of vertices and the subsets $K_1, \dots, K_N$ are called the vertex sets. Further, we call

$$E := \{(i, n_i) : i \in \{1, \dots, N\}, n_i \in \{1, \dots, L_i\}\}$$

the set of edges and we use the following notations:

$$p_e := p_{in} \text{ and } w_e := w_{in} \text{ for } e := (i, n) \in E.$$

Each edge is provided with a direction (an arrow) by marking an initial vertex through the map

$$\begin{aligned} i : E &\longrightarrow V \\ (j, n) &\longmapsto j. \end{aligned}$$

The terminal vertex $t(j, n) \in V$ of an edge $(j, n) \in E$ is determined by the corresponding map through

$$t((j, n)) := k : \Longleftrightarrow w_{jn}(K_j) \subset K_k.$$

We call the quadruple $G := (V, E, i, t)$ a directed (multi)graph or digraph. A sequence (finite or infinite) $(\dots, e_{-1}, e_0, e_1, \dots)$ of edges which corresponds to a walk along the arrows of the digraph (i.e. $t(e_k) = i(e_{k+1})$) is called a path.

Definition 2 We call the family $\mathcal{M} := (K_{i(e)}, w_e, p_e)_{e \in E}$ a (finite) Markov system.

The definition can be easily generalized to the infinite case.

The Markov system defines a random dynamical system on K by extending the probability functions $p_e|_{K_{i(e)}}$ on the whole space by zero and the maps arbitrarily, as in Remark 1.

Definition 3 We call a Markov system *irreducible* or *aperiodic* if and only if its directed graph is irreducible or aperiodic respectively.

Definition 4 (CMS) We call Markov system $\mathcal{M}$ *contractive* with an *average contracting rate* $0 < a < 1$ if and only if it satisfies the following *condition of contractiveness on average*:

$$\sum_{e \in E} p_e(x) d(w_e(x), w_e(y)) \leq a d(x, y) \text{ for all } x, y \in K_i \text{ and } i \in \{1, \dots, N\} \quad (1)$$

(it is understood here that p_e 's are extended on the whole space by zero and w_e 's arbitrarily). This condition was discovered by Richard Isaac in 1962 for the case $N = 1$ [18].

It was shown in [35] that an irreducible contractive Markov system $\mathcal{M}$ with uniformly continuous probabilities $p_e|_{K_{i(e)}} > 0$ has a unique invariant Borel probability measure if $P_x \ll P_y$ for all $x, y \in K_{i(e)}$, $e \in E$, and the subsets K_i form an open partition of K (this was shown in [35] for some locally compact spaces, but it holds also on complete separable spaces, as contractive $\mathcal{M}$ also posses invariant measures on such spaces [15]).

1.3 Fundamental Markov systems

Now, we intend to show that with every random dynamical system $\mathcal{D}$ is associated an equivalent Markov system $\mathcal{M}' := (K'_{i(e)}, w'_e, p'_e)_{e \in E'}$ (not necessarily finite) such that $P'_x \ll P'_y$ for all $x, y \in K'_{i(e)}$, $e \in E'$, and each $K'_{i(e)}$ is the largest with such property, where P'_x are the probability measures on the code space of $\mathcal{M}'$.

The construction of $\mathcal{M}'$ goes as follows. Define an equivalence relation between $x, y \in K$ by

$$x \sim y \quad :\Leftrightarrow \quad P_x \ll\!\!\gg P_y,$$

where $P_x \ll\!\!\gg P_y$ means P_x is absolutely continuous with respect to P_y and P_y is absolutely continuous with respect to P_x . Let

$$\bigsqcup_{i \in V'} K'_i = K$$

be the partition of K into the equivalence classes. Then, for every $e \in E$ and $x, y \in K'_i$, $i \in V'$,

$$p_e(x) = 0 \quad \Leftrightarrow \quad p_e(y) = 0.$$

Hence, for every $e \in E$ and $i \in V'$,

$$\text{either } p_e|_{K'_i} = 0 \text{ or } p_e|_{K'_i} > 0. \quad (2)$$

Furthermore, holds the following.

Proposition 1 *For every $e \in E$ and $i \in V'$ with $p_e|_{K'_i} > 0$, there exists $j \in V'$ such that $w_e(K'_i) \subset K'_j$.*

Proof. Let $x, y \in K'_i$. Observe that

$$P_x(1[e, \sigma_1, \dots, \sigma_n]) = p_e(x)P_{w_e(x)}(1[\sigma_1, \dots, \sigma_n])$$

for every cylinder set $1[\sigma_1, \dots, \sigma_n]$. Hence,

$$P_{w_e(x)}(B) = \frac{P_x(S^{-1}(B) \cap 1[e])}{p_e(x)}$$

for every Borel $B \subset \Sigma^+$. Since the analogous formula holds true also for $P_{w_e(y)}$, we conclude that $w_e(x) \sim w_e(y)$. Thus, there exists $j \in V'$ such that $w_e(K'_i) \subset K'_j$. $\square$

By (2) and Proposition 1, we can define a Markov systems associated with $\mathcal{D}$.

Definition 5 Let

$$E'_i := \{(i, e) : p_e|_{K'_i} > 0, e \in E\} \quad \text{for all } i \in V'$$

and

$$E' := \bigcup_{i \in V'} E'_i.$$

For every $(i, e) \in E'$ set $p'_{(i,e)} := p_e|_{K'_i}$, $w'_{(i,e)} := w_e|_{K'_i}$, $i'((i, e)) = i$ and $t'((i, e)) = j$ where $w_e(K'_i) \subset K'_j$. Then $G' := (V', E', i', t')$ is a directed graph and we call $\mathcal{M}' := (K'_{i(e)}, w'_e, p'_e)_{e \in E'}$ the *fundamental Markov systems* associated with the random dynamical system $\mathcal{D}$.

Now, we need to show that the vertex sets of the fundamental Markov system $\mathcal{M}'$ associated with $\mathcal{D}'$ are measurable. Otherwise, possible Banach-Tarski effects might make our construction scientifically irrelevant. For that, we need to make clear the constructive nature of the equivalence relation which defines the vertex sets.

For $x, y \in K$, let

$$X_n(\sigma) := \begin{cases} \frac{P_x(1[\sigma_1, \dots, \sigma_n])}{P_y(1[\sigma_1, \dots, \sigma_n])}, & P_y(1[\sigma_1, \dots, \sigma_n]) > 0 \\ 0, & P_x(1[\sigma_1, \dots, \sigma_n]) = 0 \\ \infty, & P_x(1[\sigma_1, \dots, \sigma_n]) > 0 \text{ and } P_y(1[\sigma_1, \dots, \sigma_n]) = 0 \end{cases}$$

and

$$Y_n(\sigma) := \begin{cases} \frac{P_y(1[\sigma_1, \dots, \sigma_n])}{P_x(1[\sigma_1, \dots, \sigma_n])}, & P_x(1[\sigma_1, \dots, \sigma_n]) > 0 \\ 0, & P_y(1[\sigma_1, \dots, \sigma_n]) = 0 \\ \infty, & P_y(1[\sigma_1, \dots, \sigma_n]) > 0 \text{ and } P_x(1[\sigma_1, \dots, \sigma_n]) = 0 \end{cases}$$

for all $\sigma \in \Sigma^+$. Define

$$\xi(x, y) := \limsup_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} P_x(X_n > M) + \limsup_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} P_y(Y_n > M).$$

Observe that each $x \mapsto P_x(\cdot | [\sigma_1, \dots, \sigma_n])$ is a Borel measurable function. Therefore, each $x \mapsto P_x(X_n > M)$ is a Borel measurable function. Hence, $x \mapsto \xi(x, y)$ is a Borel measurable function for all $y \in K$. By the symmetry, also $y \mapsto \xi(x, y)$ is a Borel measurable function for all $x \in K$.

Lemma 1 *For all $x, y \in K$,*

$$\xi(x, y) = 0 \quad \text{if and only if} \quad P_x \ll\!\!\ll P_y.$$

Proof. Let $x, y \in K$. Let $\mathcal{A}_n$ be the finite σ -algebra on Σ^+ generated by the cylinders $[\sigma_1, \dots, \sigma_n]$. Now, observe that, for all $m \leq n$ and $C_m \in \mathcal{A}_m$,

$$\int_{C_m} X_n dP_y = \sum_{C_n \subset C_m} P_x(C_n) = P_x(C_m) = \int_{C_m} X_m dP_y. \quad (3)$$

Hence, $(X_n, \mathcal{A}_n)_{n \in \mathbb{N}}$ is a P_y -martingale. Analogously, $(Y_n, \mathcal{A}_n)_{n \in \mathbb{N}}$ is a P_x -martingale. Moreover, by (3),

$$P_x(X_n > M) = \int_{\{X_n > M\}} X_n dP_y$$

and analogously

$$P_y(Y_n > M) = \int_{\{Y_n > M\}} Y_n dP_x.$$

Hence,

$$\xi(x, y) = \limsup_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} \int_{\{X_n > M\}} X_n dP_y + \limsup_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} \int_{\{Y_n > M\}} Y_n dP_x.$$

Therefore, $\xi(x, y) = 0$ if and only if X_n and Y_n are uniformly integrable martingales. Hence, the condition $\xi(x, y) = 0$ implies that there exists $X \in \mathcal{L}^1(P_y)$ and $Y \in \mathcal{L}^1(P_x)$ such that $X_n \rightarrow X$ and $Y_n \rightarrow Y$ both in $\mathcal{L}^1$ sense, and $E_{P_y}(X | \mathcal{A}_m) = X_m$ P_y -a.e. and $E_{P_x}(Y | \mathcal{A}_m) = Y_m$ P_x -a.e. for all m . Then, by (3),

$$\int_{C_m} X dP_y = \int_{C_m} X_m dP_y = P_x(C_m) \text{ for all } C_m \in \mathcal{A}_m.$$

Hence, the Borel probability measures $X P_y$ and P_x agree on all cylinder subsets of Σ^+ , and therefore, are equal. Analogously, $Y P_x = P_y$. Thus, $P_x \ll\!\!\ll P_y$.

Conversely, $P_x \ll\!\!\ll P_y$ implies that X_n and Y_n are uniformly integrable [7], i.e. $\xi(x, y) = 0$. $\square$

Remark 2 Note that it is not obvious from the definition of ξ that the relation $\xi(x, y) = 0$ is transitive.

Proposition 2 (i) *The vertex sets K'_i , $i \in V'$, are Borel measurable.*
(ii) *Consider all probability functions $p'_e|_{K'_{i(e)}}$, $e \in E'$, to be extended on K by zero and all maps $w'_e|_{K'_{i(e)}}$, $e \in E'$, to be extended on K arbitrarily. Let U' be the Markov operator associated with the Markov system $\mathcal{M}'$. Then $U' = U$, i.e. $\mathcal{M}'$ is an equivalent random dynamical system to $\mathcal{D}$.*

Proof. (i) Let $i \in V'$. Fix $y \in K'_i$ and set $f(x) := \xi(x, y)$ for all $x \in K$. Then, by Lemma 1, $K'_i = f^{-1}(\{0\})$. Hence, as f is Borel measurable, K'_i is Borel measurable.

(ii) Let g be a bounded Borel measurable function on K and $x \in K$. Then there exists a unique $i \in V'$ such that $x \in K'_i$. Hence, by the definition of $\mathcal{M}$,

$$\begin{aligned} U'g(x) &= \sum_{e \in E'} p'_e(x)g \circ w'_e(x) = \sum_{e \in E'_i} p'_e(x)g \circ w'_e(x) = \sum_{e \in E} p_e(x)g \circ w_e(x) \\ &= Ug(x). \end{aligned}$$

□

Example 1 Suppose the random dynamical system $\mathcal{D}$ is given by the contractive Markov system $\mathcal{M}$ such that the vertex sets $K_1, \dots, K_N$ form an open partition of K and the probability functions $p_e|_{K_{i(e)}}$ are bounded away from zero and have a square summable variation, i.e. $\sum_{n \in \mathbb{N}} \phi^2(a^n) < \infty$, where ϕ is the maximum of modules of uniform continuity of functions $p_e|_{K_{i(e)}}$, $e \in E$. Then, by Lemma 2 in [35], $P_x \ll P_y$ for all $x, y \in K_i$, $i = 1, \dots, N$ (note that the openness of the partition was required in [35] only to insure that $\mathcal{M}$ has an invariant measure (Feller property)). Therefore, the fundamental Markov system associated with $\mathcal{M}$ is $\mathcal{M}$ itself.

Example 2 Let $\mathcal{D}_2 := ([0, 1], w_e, p_e)_{e=0,1}$ be the random dynamical system where $w_0(x) = x/3$, $w_1(x) = x/3 + 1/3$ for all $x \in [0, 1]$,

$$p_0(x) = \begin{cases} 0, & 0 \leq x \leq \frac{1}{9} \\ b, & \frac{1}{9} < x \leq 1 \end{cases},$$

with $0 < b < 1$, and $p_1 = 1 - p_0$.

Claim 1 *The vertex sets of the fundamental Markov system associated with $\mathcal{D}_2$ are $K_0 := [0, 1/9]$, $K_1 := (1/9, 1/3]$ and $K_2 := (1/3, 1]$.*

Proof. First, observe that partition $K_0 \cup K_1 \cup K_2$ makes a Markov system with constant probabilities from $\mathcal{D}_2$, where the transition matrix associated with it

is

$$A_2 := \begin{pmatrix} 0 & 0 & 1 \\ b & 0 & 1-b \\ 0 & b & 1-b \end{pmatrix}.$$

Therefore,

$$p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_1} x) = p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_1} y)$$

for all $x, y \in K_i$, $i = 0, 1, 2$ and $e_1, \dots, e_n \in \{0, 1\}$. Thus

$$P_x = P_y \text{ for all } x, y \in K_0, K_1, K_2.$$

It remains to show that K_0 , K_1 and K_2 are the largest with the property that $P_x <<>> P_y$ for all $x, y \in K_i$, $i = 0, 1, 2$. Let $x \in K_0$ and $y \notin K_0$. Then $P_x(1[0]) = 0$, but $P_y(1[0]) = b > 0$. Now, let $x \in K_1$ and $y \notin K_1$. If $y \in K_0$, then $P_y(1[0]) = 0$, but $P_x(1[0]) = b$. Otherwise, if $y \in K_2$, $P_x(1[00]) = 0$, but $P_y(1[00]) = b$. The claim follows. $\square$

Now, we can apply Theorem 2 in [36] to an equivalent fundamental Markov system on a disconnected set, the vertex sets of which are $\tilde{K}_0 := [0, 1/9]$, $\tilde{K}_1 := [2/9, 1/3]$ and $\tilde{K}_2 := [2/3, 1]$. (Note that there is a missprint in [36] on page 471. It should be $\tilde{A} := D^{-1}A^tD$.) By Theorem 2 in [36], $\mathcal{D}_2$ has a unique invariant Borel probability measure μ_2 with $\mu_2(K_0) = b^2/(1+b+b^2)$, $\mu_2(K_1) = b/(1+b+b^2)$ and $\mu_2(K_2) = 1/(1+b+b^2)$ and the Markov chain associated with $\mathcal{D}_2$ is geometrically ergodic with a relative rate of convergence in Monge-Kantorovich metric less or equal to $\max\{1/3, b\}^{1/2}$.

If we replace p_0 with

$$p_0(x) = \begin{cases} 0, & 0 \leq x \leq \frac{1}{27} \\ b, & \frac{1}{27} < x \leq 1 \end{cases},$$

the same argumentation as in the proof of Claim 1 shows that the fundamental Markov system associated with modified $\mathcal{D}_2$ has four vertex sets $K_0 := [0, 1/27]$, $K_1 := (1/27, 1/9]$, $K_2 := (1/9, 1/3]$ and $K_3 := (1/3, 1]$ with constant probabilities on them given by the transition Matrix

$$A_3 := \begin{pmatrix} 0 & 0 & 0 & 1 \\ b & 0 & 0 & 1-b \\ 0 & b & 0 & 1-b \\ 0 & 0 & b & 1-b \end{pmatrix}.$$

Analogously, applying Theorem 2 in [36] gives that the modified $\mathcal{D}_2$ has an attractive invariant probability measure μ_3 with the weights on vertex sets $\mu_3(K_0) = b^3/(1+b+b^2+b^3)$, $\mu_3(K_1) = b^2/(1+b+b^2+b^3)$, $\mu_3(K_2) = b/(1+b+b^2+b^3)$ and $\mu_3(K_3) = 1/(1+b+b^2+b^3)$. (Somewhat surprisingly, the estimation of the relative rate of convergence by Theorem 2 in [36] remains the same (the absolute value of eigenvalues of such transition matrices remains the same and equals b). Note that in case of $p_0 = b$ the rate of convergence to the stationary state in the Monge-Kantorovich metric is not greater than $1/3$.)

Example 3 Let $\mathcal{D}_3 := ([0, 1], w_e, p_e)_{e=0,1}$ be the random dynamical system where $w_0(x) = x/3$, $w_1(x) = x/3 + 1/3$ for all $x \in [0, 1]$,

$$p_0(x) = \begin{cases} \frac{1}{4}, & 0 \leq x \leq \frac{1}{2} \\ \frac{1}{3}, & \frac{1}{2} < x \leq 1 \end{cases}$$

and $p_1 = 1 - p_0$. As $p_{\sigma_2}(w_{\sigma_1}x) = p_{\sigma_2}(w_{\sigma_1}x)$ for all $x, y \in [0, 1]$, $\sigma_1, \sigma_2 \in \{0, 1\}$ and $p_0 > 0$, follows $P_x << P_y$ for all $x, y \in [0, 1]$. Thus the fundamental Markov system associated with $\mathcal{D}_3$ has the single vertex set.

Example 4 Let $\mathcal{D}_4 := (\mathbb{R}, w_e, p_e)_{e=0,1}$ be the random dynamical system where $w_0(x) = x/2$, $w_1(x) = x/2 + 1/2$ for all $x \in \mathbb{R}$,

$$p_0(x) = \begin{cases} \frac{1}{4}, & x \in \mathbb{Q} \\ \frac{1}{3}, & x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$

and $p_1 = 1 - p_0$. If x is rational, then all images of x under w_0 and w_1 are rational. Therefore, P_x is the Bernoulli measure generated with probabilities $\{1/4, 3/4\}$. Analogously, for irrational y , P_y is the Bernoulli measure generated with probabilities $\{1/3, 2/3\}$. Since P_x and P_y can not be absolutely continuous (absolutely continuous ergodic measures are equal), the fundamental Markov system associated with $\mathcal{D}_4$ has two vertex sets $K_0 := \mathbb{Q}$ and $K_1 := \mathbb{R} \setminus \mathbb{Q}$.

Having obtained the well defined fundamental Markov system associated with $\mathcal{D}$, we can define a Borel probability measure P'_x on $\Sigma'^+ := \{(\sigma_1, \sigma_2, \dots) : \sigma_i \in E' \forall i \in \mathbb{N}\}$ (provided with the product topology of discrete topologies) by

$$P'_x(1[e_1, \dots, e_n]) := p'_{e_1}(x) \dots p'_{e_n}(w'_{e_{n-1}} \circ \dots \circ w'_{e_1}x)$$

for all cylinder sets $1[e_1, \dots, e_n] \subset \Sigma'^+$, for every $x \in K$.

Now, we need to establish a relation between measures P'_x and P_x .

Since for each $e' \in E'$, $w'_{e'}|_{K'_{i(e')}} is a restriction of a unique w_e , we can define a map $\psi : E' \rightarrow E$ by $\psi(e') = e$. This gives a Borel-Borel-measurable map $\Psi : \Sigma'^+ \rightarrow \Sigma^+$ by $(\Psi(\sigma))_i := \psi(\sigma_i)$ for all $i \in \mathbb{N}$ and for all $\sigma \in \Sigma'^+$.$

Lemma 2 For every $x \in K$, $P_x = \Psi(P'_x)$.

Proof. Let $1[e_1, \dots, e_n] \subset \Sigma^+$ be a cylinder set. Then, by the definition of Ψ , we can write $\Psi^{-1}(1[e_1, \dots, e_n])$ as a disjoint union of some cylinder sets $1[e'_1, \dots, e'_n] \subset \Sigma'^+$, i.e.

$$\Psi^{-1}(1[e_1, \dots, e_n]) = \bigcup_{(e'_1, \dots, e'_n), \psi(e'_i) = e_i} 1[e'_1, \dots, e'_n].$$

Therefore,

$$\Psi(P'_x)(1[e_1, \dots, e_n]) = \sum_{(e'_1, \dots, e'_n), \psi(e'_i) = e_i} p'_{e'_1}(x) \dots p'_{e'_n}(w'_{e'_{n-1}} \circ \dots \circ w'_{e'_1}x).$$

Now, observe that, by the definition of p'_e ,

$$\sum_{e', \psi(e')=e} p'_{e'} = p_e.$$

This implies that

$$\Psi(P'_x)(1[e_1, \dots, e_n]) = p_{e_1}(x) \dots p_{e_n} \circ w_{e_{n-1}} \circ \dots \circ w_{e_1}(x) = P_x(1[e_1, \dots, e_n]).$$

Thus, the claim follows. $\square$

Proposition 3 *For all $x, y \in K$,*

$$P_x \ll P_y \iff P'_x \ll P'_y.$$

Proof. $' \Leftarrow'$ Let $A \subset \Sigma^+$ Borel measurable such that $P_y(A) = 0$. Then, by Lemma 2, $P'_y(\Psi^{-1}(A)) = 0$. Hence, by Lemma 2 and the hypothesis, $P_x(A) = P'_x(\Psi^{-1}(A)) = 0$. Thus $P_x \ll P_y$.

$' \Rightarrow'$ First, observe that, by the construction of M' , for every cylinder set $1[e'_1, \dots, e'_n] \subset \Sigma'^+$ with $P'_y(1[e'_1, \dots, e'_n]) > 0$, there exists a unique cylinder set $1[e_1, \dots, e_n] \subset \Sigma^+$ such that $P'_y(1[e'_1, \dots, e'_n]) = P_y(1[e_1, \dots, e_n])$ and $\Psi(1[e'_1, \dots, e'_n]) = 1[e_1, \dots, e_n]$. Hence

$$P'_y(1[e'_1, \dots, e'_n]) = P_y(\Psi(1[e'_1, \dots, e'_n])). \quad (4)$$

Now, let $B \subset \Sigma'$ Borel measurable such that $P'_y(B) = 0$. Let $\epsilon > 0$. By the hypothesis, there exists $\delta > 0$ such that

$$P_y(C) < \delta \implies P_x(C) < \epsilon \quad (5)$$

for all Borel measurable $C \subset \Sigma^+$. By the Borel regularity of P'_x , there exists a countable family of cylinder sets $C_k \subset \Sigma'^+$, $k \in \mathbb{N}$, such that $B \subset \bigcup_k C_k$ and $P'_y(\bigcup_k C_k) < \delta$. Since we can write every finite union $\bigcup_k^m C_k$ as a disjoint union of cylinder sets $\bigcup_k^{n_m} \tilde{C}_k$,

$$\sum_{k=1}^{\infty} P'_y(\tilde{C}) = P'_y(\bigcup_k C_k) < \delta.$$

Hence, by (4) and the Lemma 2,

$$P_y(\bigcup_k \Psi(C_k)) \leq \sum_{k=1}^{\infty} P_y(\Psi(\tilde{C})) = \sum_{k=1}^{\infty} P'_y(\tilde{C}) < \delta.$$

Therefore, by (5) and Lemma 2,

$$P'_x(B) \leq P'_x(\Psi^{-1}(\Psi(B))) = P_x(\Psi(B)) \leq P_x(\bigcup_k \Psi(\tilde{C}_k)) < \epsilon.$$

Since ϵ was arbitrary, this completes the proof. $\square$

Theorem 1 *Suppose $\mathcal{D}$ is a random dynamical system with finitely many uniformly continuous probability functions p_e and continuous maps w_e on a complete separable metric space K . Suppose that the fundamental Markov system associated with $\mathcal{D}$ has finitely many vertices, is irreducible and contractive. Then*

- (i) $\mathcal{D}$ has a unique invariant Borel probability measure μ .
- (ii) For every $x \in K$,

$$\frac{1}{n} \sum_{k=0}^{n-1} f \circ w_{\sigma_k} \circ \dots \circ w_{\sigma_1}(x) \rightarrow \int f d\mu \text{ for } P_x\text{-a.e. } \sigma \in \Sigma^+$$

for all bounded continuous functions f .

Proof. Apply Theorem 4 in [35] for the fundamental Markov system associated with $\mathcal{D}$ with the following justifications. In [35], the vertex sets were required to form an open partition of a state space in which sets of finite diameter are relatively compact. This was to insure that the Markov operator has the Feller property and an invariant Borel probability measure. Here, the Feller property is already given by $\mathcal{D}$ and the existence of invariant measures for our fundamental Markov system associated with $\mathcal{D}$ (on complete separable space) was shown in [15]. Also, it was required in [35] that the probability functions $p_e|_{K_{i(e)}}$ shall be bounded away from zero, but it was only required for the proof that $P_x \ll P_y$ for all x and y in the same vertex set (Lemma 2 in [35]). The latter is given here by the construction of the fundamental Markov system associated with $\mathcal{D}$ and Proposition 3.

After obtaining the result for the fundamental Markov system associated with $\mathcal{D}$, deduce the result for $\mathcal{D}$ by Proposition 3. $\square$

Remark 3 Note that the fundamental Markov system $\mathcal{M}'$ associated with $\mathcal{D}$ is contractive if $\mathcal{D}$ is contractive.

Remark 4 Recall that there exist contractive random dynamical systems with strictly positive continuous probability functions which have more than one probability measure [8],[6]. By Theorem 1 the fundamental Markov system associated with such a random dynamical system can not have a single vertex set.

Conjecture 1 I believe that fundamental Markov systems resolve the question of the necessary and sufficient condition for the stability of such random dynamical systems, which has been open already for more than 70 years, in the following way. *The random dynamical system has a unique invariant Borel probability measure if and only if the fundamental Markov system associated with it is recurrent (every vertex of it is reached from any other by a finite path).* Note that a recurrent Markov system is necessarily countable (every vertex of it can be coded by a finite path).

References

- [1] M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo, Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **24** (1988) 367-394.
- [2] M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo, Erratum: Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **25** (1989) 589-590.
- [3] M. F. Barnsley, J. H. Elton and D. P. Hardin, Recurrent iterated function systems, *Constructive Approximation* **5** (1989) 3-31.
- [4] M. F. Barnsley, A. Deliu, R. Xie, Stationary stochastic processes and fractal data compression, *Int. J. Bifurcation & Chaos* **7** (1997), 551-567.
- [5] H. Berbee, Chains with Infinite Connections: Uniqueness and Markov Representation, *Probab. Th. Rel. Fields* **76** (1987), 243-253.
- [6] N. Berger, Ch. Hoffman, V. Sidoravicius, Nonuniqueness for specifications in $\ell^{2+\epsilon}$, *arXiv:math/0312344v4*.
- [7] P. Billingsley, *Probability and Measure*, John Wiley and Sons (1979).
- [8] M. Bramson, S. Kalikow *Nonuniqueness in g-functions*, *Israel J. Math.* **84** (1993) 153-160.
- [9] L. Breiman, The strong law of large numbers for a class of Markov chains, *Ann. Math. Stat.* **31** (1960) 801-803.
- [10] P. Brémaud, *Markov Chains: Gibbs fields, Monte Carlo simulation, and queues*, Springer (1998).
- [11] G. P. Basharina, A. N. Langville and V. A. Naumov, The life and work of A.A. Markov, *Linear Algebra and its Applications* **386** (2004) 3-26.
- [12] W. Doeblin and R. Fortet, Sur les chaînes à liaisons complètes, *Bull. Soc. Math. France* **65** (1937) 132-148.
- [13] J. H. Elton, An ergodic theorem for iterated maps, *Ergod. Th. & Dynam. Sys.* **7** (1987) 481-488.
- [14] J. H. Elton and Z. Yan, Approximation of measures by Markov processes and homogeneous affine iterated function systems, *Constructive Approximation* **5** (1989) 69-87.
- [15] K. Horbacz and T. Szarek, Irreducible Markov systems on Polish spaces, *Studia Math.* **177** (2006), 285-295.
- [16] J. Hutchinson, Fractals and Self-Similarity, *Indiana U. J. of Math.* **30** (1981), 713-747.

- [17] M. Iosifescu, Ș. Grigorescu, *Dependence with complete connections and its applications*, Cambridge Tracts in Mathematics, 96. Cambridge University Press, Cambridge, (1990).
- [18] R. Isaac, Markov processes and unique stationary probability measures, *Pacific J. Math.* **12** (1962) 273-286.
- [19] A. Johansson and A. Öberg, Square summability of variations of g -functions and uniqueness of g -measures, *Math. Res. Lett.* **10** (2003) 587-601.
- [20] S. Karlin, Some random walks occurring in learning models, *Pacific J. Math.* **3** (1953) 725-756.
- [21] M. Keane, Strongly Mixing g -Measures, *Inventiones math.* **16** (1972) 309-324.
- [22] T. Kaijser, On a new contraction condition for random systems with complete connections, *Rev. Roumaine Math. Pures Appl.* **26** (1981) 1075-1117.
- [23] Y. Kifer and P.-D. Liu, *Random dynamics*, Handbook of Dynamical Systems, vol. **1B** (B. Hasselblatt and A. Katok—eds.), 379-499, North-Holland/Elsevier (2006).
- [24] A. Kwiecińska and W. Ślomyński, Random dynamical systems arising from iterated function systems with place-dependent probabilities, *Statistics & Probability Letters* **50** (2000) 401-407.
- [25] F. Ledrappier, Principe variationnel et systèmes dynamiques symboliques, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **30** (1974) 185-202.
- [26] A. A. Markov, Extension of the law of large numbers to dependent events (in Russian), *Bull. Soc. Phys. Math. Kazan*, **2** (1906), no. 15, 155-156.
- [27] O. Onicescu and G. Mihoc, Sur les chaînes de variables statistiques, *Bull. Sci. Math. de France* **59** (1935) 174-192.
- [28] D. S. Ornstein, Bernoulli shifts with the same entropy are isomorphic, *Adv. in Math.* **4** (1970), 339-348.
- [29] D. S. Ornstein and P. Shields, An uncountable family of K -automorphisms, *Adv. in Math.* **10** (1973), 63-88.
- [30] Ö. Stenflo, Uniqueness in g -measures, *Nonlinearity* **16** (2003) 403-410.
- [31] W. Ślomyński, *Dynamical entropy, Markov operators, and iterated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).
- [32] T. Szarek, Invariant measures for nonexpansive Markov operators on Polish spaces, *Diss. Math.* **415**, 1-62 (2003).
- [33] P. Walters, Ruelle's Operator Theorem and g -measures, *Tran. AMS* **214** (1975) 375-387.

- [34] I. Werner, Contractive Markov systems, *J. London Math. Soc.* **71** (2005) 236-258.
- [35] I. Werner, Contractive Markov systems II, *arXiv:math/0506476*.
- [36] I. Werner, Contractive Markov system with constant probabilities, *J. Theoret. Prob.* **18** (2005), no. 2, 469-479.
- [37] I. Werner, A necessary condition for the uniqueness of the stationary state of a Markov system, *arXiv:math/0508054*.
- [38] I. Werner, On coding with Feller contractive Markov systems, *arXiv:math/0506476*.